arXiv:0904.1115v1 [math.CA] 7 Apr 2009

THE FUNCTION $(b^x - a^x)/x$: RATIO'S PROPERTIES

BAI-NI GUO AND FENG QI

Abstract. In the paper, after reviewing the history, background, origin, and applications of the functions $\frac{b^t-a^t}{t}$ and $\frac{e^{-\alpha t}-e^{-\beta t}}{1-e^{-t}}$, we establish sufficient and necessary conditions such that the special function $\frac{e^{\alpha t}-e^{\beta t}}{e^{\lambda t}-e^{\mu t}}$ are monotonic, logarithmic convex, logarithmic concave, 3-log-convex and 3-log-concave on $\mathbb{R}$, where α,β,λ and μ are real numbers satisfying $(\alpha,\beta)\ne(\lambda,\mu)$, $(\alpha,\beta)\ne(\mu,\lambda)$, $\alpha\ne\beta$ and $\lambda\ne\mu$.

1. Introduction

Recall [42, 43] that a k-times differentiable function $f(t)>0$ is said to be k-log-convex on an interval I if

$$0\le[\ln f(t)]^{(k)}<\infty,\quad k\in\mathbb{N} \tag{1}$$

on I; If the inequality (1) reverses then f is said to be k-log-concave on I.

For $b>a>0$, let

$$G_{a,b}(t)=\begin{cases}\dfrac{b^t-a^t}{t}, & t\ne0;\\ \ln b-\ln a, & t=0.\end{cases} \tag{2}$$

In [47, 48], the complete monotonicity and inequality properties of $G_{a,b}(t)$ were first investigated. In [9, 29, 37, 38], the 3-log-convex and 3-log-concave properties of $G_{a,b}(t)$ were shown. The function $G_{a,b}(t)$ has close relationships with the incomplete gamma function [20, 26, 27]. It was ever used to prove the Schur-convex properties [20, 26, 38], the logarithmic convexities [3, 4, 23, 24, 29, 38], the monotonicity [44, 49] of the extended mean values. It was applied in [7, 30, 31, 32, 35, 36] to construct Steffensen pairs. It was also employed in [41] to verify Elezović-Giordano-Pečarić's theorem [6, Theorem 1] which is related to the monotonicity of a function involving the ratio of two gamma functions. For more information, please refer to [1, 27] and closely-related references therein.

For $b>a>0$, let

$$F_{a,b}(t)=\begin{cases}\dfrac{t}{e^{bt}-e^{at}}, & t\ne0;\\ \dfrac{1}{b-a}, & t=0.\end{cases} \tag{3}$$

In [2, 8, 14, 19, 50, 51, 52], [5, p. 217] and [13, p. 295], the inequalities, monotonicity and logarithmic convexities of the function $F_{a,b}(t)$ for $a=b-1$ and its logarithmic derivatives of the first and second orders are established. In [14], the history, background and origin of $F_{a,b}(t)$ for $a=b-1$ and its first two logarithmic derivatives were cultivated. In [12, 45, 46], the logarithmic derivative of $F_{a,b}(t)$ for $a=b-1$ was applied to study the complete monotonicity of remainders of the first Binet formula

---

2000 *Mathematics Subject Classification.* Primary 33B10; Secondary 26A48.

Key words and phrases. Monotonicity, logarithmic convexity, logarithmic concavity, 3-log-convex function, 3-log-concave function, exponential function.

The second author was partially supported by the China Scholarship Council.

This paper was typeset using $\mathcal{A}_{\mathcal{M}}\mathcal{S}$-LaTeX.

and the psi function. In [10, 15, 16, 17, 34], the function $F_{\ln a,\ln b}(t)$ was utilized to generalize Bernoulli numbers and polynomials. In [9, 37], the 3-log-convex and 3-log-concave properties of $F_{a,b}(t)$ were shown, among other things.

For real numbers α and β satisfying $\alpha\ne\beta$, $(\alpha,\beta)\ne(0,1)$ and $(\alpha,\beta)\ne(1,0)$, let

$$Q_{\alpha,\beta}(t)=\begin{cases}\dfrac{e^{-\alpha t}-e^{-\beta t}}{1-e^{-t}}, & t\ne0;\\ \beta-\alpha, & t=0.\end{cases} \tag{4}$$

In [11, 25, 28, 41], the monotonicity and logarithmic convexities of $Q_{\alpha,\beta}(t)$ were discussed and the following conclusions were procured:

1. The function $Q_{\alpha,\beta}(t)$ is increasing on $(0,\infty)$ if and only if $(\beta-\alpha)(1-\alpha-\beta)\ge0$ and $(\beta-\alpha)(|\alpha-\beta|-\alpha-\beta)\ge0$;
2. The function $Q_{\alpha,\beta}(t)$ is decreasing on $(0,\infty)$ if and only if $(\beta-\alpha)(1-\alpha-\beta)\le0$ and $(\beta-\alpha)(|\alpha-\beta|-\alpha-\beta)\le0$;
3. The function $Q_{\alpha,\beta}(t)$ is increasing on $(-\infty,0)$ if and only if $(\beta-\alpha)(1-\alpha-\beta)\ge0$ and $(\beta-\alpha)(2-|\alpha-\beta|-\alpha-\beta)\ge0$;
4. The function $Q_{\alpha,\beta}(t)$ is decreasing on $(-\infty,0)$ if and only if $(\beta-\alpha)(1-\alpha-\beta)\le0$ and $(\beta-\alpha)(2-|\alpha-\beta|-\alpha-\beta)\le0$;
5. The function $Q_{\alpha,\beta}(t)$ is increasing on $(-\infty,\infty)$ if and only if $(\beta-\alpha)(|\alpha-\beta|-\alpha-\beta)\ge0$ and $(\beta-\alpha)(2-|\alpha-\beta|-\alpha-\beta)\ge0$;
6. The function $Q_{\alpha,\beta}(t)$ is decreasing on $(-\infty,\infty)$ if and only if $(\beta-\alpha)(|\alpha-\beta|-\alpha-\beta)\le0$ and $(\beta-\alpha)(2-|\alpha-\beta|-\alpha-\beta)\le0$;
7. The function $Q_{\alpha,\beta}(t)$ on $(-\infty,\infty)$ is logarithmically convex if $\beta-\alpha>1$ and logarithmically concave if $0<\beta-\alpha<1$;
8. If $1>\beta-\alpha>0$, then $Q_{\alpha,\beta}(t)$ is 3-log-convex on $(0,\infty)$ and 3-log-concave on $(-\infty,0)$; if $\beta-\alpha>1$, then $Q_{\alpha,\beta}(t)$ is 3-log-concave on $(0,\infty)$ and 3-log-convex on $(-\infty,0)$.

The monotonicity of $Q_{\alpha,\beta}(t)$ on $(0,\infty)$ was applied in [11, 18, 39, 40] to present necessary and sufficient conditions such that some functions involving ratios of the gamma and q-gamma functions are logarithmically completely monotonic. The logarithmic convexities of $Q_{\alpha,\beta}(t)$ on $(0,\infty)$ was used in [33, 41] to supply alternative proofs for Elezović-Giordano-Pečarić's theorem. For detailed information, please refer to [21, 22] and related references therein.

The functions $G_{a,b}(t)$, $F_{a,b}(t)$ and $Q_{\alpha,\beta}(t)$ have the following relations:

$$\begin{aligned} G_{a,b}(t)&=\frac{1}{F_{\ln a,\ln b}(t)}, & F_{a,b}(t)&=\frac{1}{G_{e^b,e^a}(t)},\\ Q_{\alpha,\beta}(t)&=\frac{G_{e^{-\alpha},e^{-\beta}}(t)}{G_{1,e^{-1}}(t)}, & Q_{\alpha,\beta}(t)&=\frac{F_{0,-1}(t)}{F_{-\alpha,-\beta}(t)}.\end{aligned} \tag{5}$$

For real numbers α,β,λ and μ satisfying $(\alpha,\beta)\ne(\lambda,\mu)$, $(\alpha,\beta)\ne(\mu,\lambda)$, $\alpha\ne\beta$ and $\lambda\ne\mu$, let

$$H_{\alpha,\beta;\lambda,\mu}(t)=\begin{cases}\dfrac{e^{\alpha t}-e^{\beta t}}{e^{\lambda t}-e^{\mu t}}, & t\ne0,\\ \dfrac{\beta-\alpha}{\lambda-\mu}, & t=0.\end{cases} \tag{6}$$

For positive numbers r,s,u and v satisfying $(r,s)\ne(u,v)$, $(r,s)\ne(v,u)$, $r\ne s$ and $u\ne v$, let

$$P_{r,s;u,v}(t)=\begin{cases}\dfrac{r^t-s^t}{u^t-v^t}, & t\ne0,\\ \dfrac{\ln r-\ln s}{\ln u-\ln v}, & t=0.\end{cases} \tag{7}$$

It is clear that

$$H_{\alpha,\beta;\lambda,\mu}(t) = P_{e^\alpha,e^\beta;e^\lambda,e^\mu}(t) \quad \text{and} \quad P_{r,s;u,v}(t) = H_{\ln r,\ln s;\ln u,\ln v}(t). \tag{8}$$

In addition, the functions $H_{\alpha,\beta;\lambda,\mu}(t)$ and $P_{r,s;u,v}(t)$ can be represented as ratios of $G_{a,b}(t)$, $F_{a,b}(t)$ and $Q_{\alpha,\beta}(t)$ as follows:

$$H_{\alpha,\beta;\lambda,\mu}(t) = \frac{F_{\lambda,\mu}(t)}{F_{\alpha,\beta}(t)} = \frac{Q_{-\alpha,-\beta}}{Q_{-\lambda,-\mu}} \quad \text{and} \quad P_{r,s;u,v}(t) = \frac{G_{r,s}(t)}{G_{u,v}(t)}. \tag{9}$$

Since the functions $G_{a,b}(t)$, $F_{a,b}(t)$ and $Q_{\alpha,\beta}(t)$ have a long history, a deep background and many applications to several areas, so we continue to study the monotonicity and logarithmic convexities of their ratios, $H_{\alpha,\beta;\lambda,\mu}(t)$ and $P_{r,s;u,v}(t)$.

Our main results may be stated as follows.

Theorem 1. *For real numbers α,β,λ and μ with $(\alpha,\beta)\ne(\lambda,\mu)$, $(\alpha,\beta)\ne(\mu,\lambda)$, $\alpha\ne\beta$ and $\lambda\ne\mu$, let*

$$\mathcal{A} = (\alpha-\beta)(\alpha+\beta-\lambda-\mu), \tag{10}$$

$$\mathcal{B} = (\alpha-\beta)(\alpha+\beta-|\alpha-\beta|-2\lambda), \tag{11}$$

$$\mathcal{C} = (\alpha-\beta)(\alpha+\beta+|\alpha-\beta|-2\lambda), \tag{12}$$

$$\mathcal{D} = (\alpha-\beta)(\alpha+\beta+|\alpha-\beta|-2\mu), \tag{13}$$

$$\mathcal{E} = (\alpha-\beta)(\alpha+\beta-|\alpha-\beta|-2\mu). \tag{14}$$

Then the function $H_{\alpha,\beta;\lambda,\mu}(t)$ has the following properties:

1. *The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is increasing on $(0,\infty)$ if and only if either $\lambda>\mu$, $\mathcal{A}\ge0$ and $\mathcal{C}\ge0$ or $\lambda<\mu$, $\mathcal{A}\le0$ and $\mathcal{B}\le0$.*
2. *The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is decreasing on $(0,\infty)$ if and only if either $\lambda<\mu$, $\mathcal{A}\ge0$ and $\mathcal{B}\ge0$ or $\lambda>\mu$, $\mathcal{A}\le0$ and $\mathcal{C}\le0$.*
3. *The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is increasing on $(-\infty,0)$ if and only if either $\lambda>\mu$, $\mathcal{A}\ge0$ and $\mathcal{E}\ge0$ or $\lambda<\mu$, $\mathcal{A}\le0$ and $\mathcal{D}\le0$.*
4. *The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is decreasing on $(-\infty,0)$ if and only if either $\lambda>\mu$, $\mathcal{A}\le0$ and $\mathcal{E}\le0$ or $\lambda<\mu$, $\mathcal{A}\ge0$ and $\mathcal{D}\ge0$.*
5. *The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is increasing on $(-\infty,\infty)$ if and only if either $\lambda>\mu$, $\mathcal{C}\ge0$ and $\mathcal{E}\ge0$ or $\lambda<\mu$, $\mathcal{B}\le0$ and $\mathcal{D}\le0$.*
6. *The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is decreasing on $(-\infty,\infty)$ if and only if either $\lambda>\mu$, $\mathcal{C}\le0$ and $\mathcal{E}\le0$ or $\lambda<\mu$, $\mathcal{B}\ge0$ and $\mathcal{D}\ge0$.*
7. *The function $H_{\alpha,\beta;\lambda,\mu}(t)$ on $(-\infty,\infty)$ is logarithmically convex if $\frac{\alpha-\beta}{\lambda-\mu}>1$ or logarithmically concave if $0<\frac{\alpha-\beta}{\lambda-\mu}<1$.*
8. *The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is 3-log-convex on $(0,\infty)$ and 3-log-concave on $(-\infty,0)$ if either $\lambda-\mu>\alpha-\beta>0$ or $\alpha-\beta<\lambda-\mu<0$; The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is 3-log-concave on $(0,\infty)$ and 3-log-convex on $(-\infty,0)$ if either $\alpha-\beta>\lambda-\mu>0$ or $\lambda-\mu<\alpha-\beta<0$.*

Theorem 2. *For positive numbers r,s,u and v with $(r,s)\ne(u,v)$, $(r,s)\ne(v,u)$, $r\ne s$ and $u\ne v$, let*

$$\mathfrak{A} = \ln\frac{rs}{uv}\ln\frac{r}{s}, \quad \mathfrak{B} = \left(\ln\frac{rs}{u^2} - \left|\ln\frac{r}{s}\right|\right)\ln\frac{r}{s}, \quad \mathfrak{C} = \left(\ln\frac{rs}{u^2} + \left|\ln\frac{r}{s}\right|\right)\ln\frac{r}{s},$$

$$\mathfrak{D} = \left(\ln\frac{rs}{v^2} + \left|\ln\frac{r}{s}\right|\right)\ln\frac{r}{s}, \quad \mathfrak{E} = \left(\ln\frac{rs}{v^2} - \left|\ln\frac{r}{s}\right|\right)\ln\frac{r}{s}.$$

Then the function $P_{r,s;u,v}(t)$ has the following properties:

1. *The function $P_{r,s;u,v}(t)$ is increasing on $(0,\infty)$ if and only if either $u>v$, $\mathfrak{A}\ge0$ and $\mathfrak{C}\ge0$ or $u<v$, $\mathfrak{A}\le0$ and $\mathfrak{B}\le0$.*

(2) *The function* $P_{r,s;u,v}(t)$ *is decreasing on* $(0,\infty)$ *if and only if either* $u<v$, $\mathfrak{A}\ge 0$ *and* $\mathfrak{B}\ge 0$ *or* $u>v$, $\mathfrak{A}\le 0$ *and* $\mathfrak{C}\le 0$.

(3) *The function* $P_{r,s;u,v}(t)$ *is increasing on* $(-\infty,0)$ *if and only if* $u>v$, $\mathfrak{A}\ge 0$ *and* $\mathfrak{E}\ge 0$, *or* $u<v$, $\mathfrak{A}\le 0$ *and* $\mathfrak{D}\le 0$.

(4) *The function* $P_{r,s;u,v}(t)$ *is decreasing on* $(-\infty,0)$ *if and only if either* $u>v$, $\mathfrak{A}\le 0$ *and* $\mathfrak{E}\le 0$ *or* $u<v$, $\mathfrak{A}\ge 0$ *and* $\mathfrak{D}\ge 0$.

(5) *The function* $P_{r,s;u,v}(t)$ *is increasing on* $(-\infty,\infty)$ *if and only if either* $u>v$, $\mathfrak{C}\ge 0$ *and* $\mathfrak{E}\ge 0$ *or* $u<v$, $\mathfrak{B}\le 0$ *and* $\mathfrak{D}\le 0$.

(6) *The function* $P_{r,s;u,v}(t)$ *is decreasing on* $(-\infty,\infty)$ *if and only if either* $u>v$, $\mathfrak{C}\le 0$ *and* $\mathfrak{E}\le 0$ *or* $u<v$, $\mathfrak{B}\ge 0$ *and* $\mathfrak{D}\ge 0$.

(7) *The function* $P_{r,s;u,v}(t)$ *on* $(-\infty,\infty)$ *is logarithmically convex if* $\frac{\ln(r/s)}{\ln(u/v)}>1$ *or logarithmically concave if* $0<\frac{\ln(r/s)}{\ln(u/v)}<1$.

(8) *The function* $P_{r,s;u,v}(t)$ *is* 3*-log-convex on* $(0,\infty)$ *and* 3*-log-concave on* $(-\infty,0)$ *if* $\frac{u}{v}>\frac{r}{s}>1$ *or* $\frac{r}{s}<\frac{u}{v}<1$; *The function* $P_{r,s;u,v}(t)$ *is* 3*-log-concave on* $(0,\infty)$ *and* 3*-log-convex on* $(-\infty,0)$ *if* $\frac{r}{s}>\frac{u}{v}>1$ *or* $\frac{u}{v}<\frac{r}{s}<1$.

Remark 1. The monotonicity of the functions $H_{\alpha,\beta;\lambda,\mu}(t)$ and $P_{r,s;u,v}(t)$ can be described by Table 1.

Remark 2. In [28, Remark 2.1] it was remarked that the function $Q_{\alpha,\beta}(t)$ can not be either 4-log-convex or 4-log-concave in either $(-\infty,0)$ or $(0,\infty)$, saying nothing of $(-\infty,\infty)$. Therefore, neither $H_{\alpha,\beta;\lambda,\mu}(t)$ nor $P_{r,s;u,v}(t)$ is either 4-log-convex or 4-log-concave on either $(-\infty,0)$ or $(0,\infty)$, saying nothing of $(-\infty,\infty)$.

2. PROOFS OF THEOREMS

Proof of Theorem 1. For $t\ne 0$, the function $H_{\alpha,\beta;\lambda,\mu}(t)$ can rewritten as

$$H_{\alpha,\beta;\lambda,\mu}(t)=\frac{e^{(\alpha-\lambda)t}-e^{(\beta-\lambda)t}}{1-e^{(\mu-\lambda)t}}=\frac{e^{-\frac{\alpha-\lambda}{\mu-\lambda}w}-e^{-\frac{\beta-\lambda}{\mu-\lambda}w}}{1-e^{-w}}=\frac{e^{-Aw}-e^{-Bw}}{1-e^{-w}},$$

where

$$A=\frac{\alpha-\lambda}{\mu-\lambda},\quad B=\frac{\beta-\lambda}{\mu-\lambda},\quad w=(\lambda-\mu)t. \tag{15}$$

Differentiating with respect to t yields

$$H'_{\alpha,\beta;\lambda,\mu}(t)=(\lambda-\mu)Q'_{A,B}(w), \tag{16}$$

$$\begin{aligned}[\ln H_{\alpha,\beta;\lambda,\mu}(t)]''&=\left[\frac{H'_{\alpha,\beta;\lambda,\mu}(t)}{H_{\alpha,\beta;\lambda,\mu}(t)}\right]'=(\lambda-\mu)\frac{\mathrm{d}}{\mathrm{d}t}\left[\frac{Q'_{A,B}(w)}{Q_{A,B}(w)}\right]\\&=(\lambda-\mu)^2\left[\frac{Q'_{A,B}(w)}{Q_{A,B}(w)}\right]'=(\lambda-\mu)^2[\ln Q_{A,B}(w)]''\end{aligned} \tag{17}$$

and

$$[\ln H_{\alpha,\beta;\lambda,\mu}(t)]'''=(\lambda-\mu)^3\left[\frac{Q'_{A,B}(w)}{Q_{A,B}(w)}\right]''=(\lambda-\mu)^3[\ln Q_{A,B}(w)]'''. \tag{18}$$

By virtue of [25, Theorem 2] (see also [11, Theorem 3.1] and [41, Lemma 1]) and the second order derivative (17), it is easy to deduce that the function $H_{\alpha,\beta;\lambda,\mu}(t)$ is logarithmically convex if $\frac{\beta-\alpha}{\mu-\lambda}>1$ and logarithmically concave if $0<\frac{\beta-\alpha}{\mu-\lambda}<1$ on $(-\infty,\infty)$.

By virtue of [28, Theorem 1.1] and the third order derivative (18), it is not difficult to obtain that

(1) if $\lambda>\mu$ and $1>\frac{\beta-\alpha}{\mu-\lambda}>0$, then $H_{\alpha,\beta;\lambda,\mu}(t)$ is 3-log-convex on $(0,\infty)$ and 3-log-concave on $(-\infty,0)$;

Table 1. Monotoncity of the functions $H_{\alpha,\beta;\lambda,\mu}(t)$ and $P_{r,s;u,v}(t)$

Intervals	Monotonicities	$\mathcal{A}$ or $\mathfrak{A}$	$\mathcal{B}$ or $\mathfrak{B}$	$\mathcal{C}$ or $\mathfrak{C}$	$\mathcal{D}$ or $\mathfrak{D}$	$\mathcal{E}$ or $\mathfrak{E}$	λ and μ or u and v
$(0,\infty)$	increasing	≥ 0		≥ 0			$\lambda > \mu$ or $u > v$
$(0,\infty)$	increasing	≤ 0	≤ 0				$\lambda < \mu$ or $u < v$
$(0,\infty)$	decreasing	≥ 0	≥ 0				$\lambda < \mu$ or $u < v$
$(0,\infty)$	decreasing	≤ 0		≤ 0			$\lambda > \mu$ or $u > v$
$(-\infty,0)$	increasing	≥ 0				≥ 0	$\lambda > \mu$ or $u > v$
$(-\infty,0)$	increasing	≤ 0			≤ 0		$\lambda < \mu$ or $u < v$
$(-\infty,0)$	decreasing	≤ 0				≤ 0	$\lambda > \mu$ or $u > v$
$(-\infty,0)$	decreasing	≥ 0			≥ 0		$\lambda < \mu$ or $u < v$
$(-\infty,\infty)$	increasing			≥ 0		≥ 0	$\lambda > \mu$ or $u > v$
$(-\infty,\infty)$	increasing		≤ 0		≤ 0		$\lambda < \mu$ or $u < v$
$(-\infty,\infty)$	decreasing			≤ 0		≤ 0	$\lambda > \mu$ or $u > v$
$(-\infty,\infty)$	decreasing		≥ 0		≥ 0		$\lambda < \mu$ or $u < v$

(2) if $\lambda > \mu$ and $\frac{\beta-\alpha}{\mu-\lambda} > 1$, then $H_{\alpha,\beta;\lambda,\mu}(t)$ is 3-log-concave on $(0,\infty)$ and 3-log-convex on $(-\infty,0)$;
(3) if $\lambda < \mu$ and $1 > \frac{\beta-\alpha}{\mu-\lambda} > 0$, then $H_{\alpha,\beta;\lambda,\mu}(t)$ is 3-log-concave on $(0,\infty)$ and 3-log-convex on $(-\infty,0)$;
(4) if $\lambda < \mu$ and $\frac{\beta-\alpha}{\mu-\lambda} > 1$, then $H_{\alpha,\beta;\lambda,\mu}(t)$ is 3-log-convex on $(0,\infty)$ and 3-log-concave on $(-\infty,0)$.

Direct computation gives

$$(B-A)(1-A-B)=\frac{\mathcal{A}}{(\lambda-\mu)^2};$$

$$(B-A)(|A-B|-A-B)=\begin{cases}\dfrac{\mathcal{B}}{(\lambda-\mu)^2}, & \lambda<\mu,\\[2mm] \dfrac{\mathcal{C}}{(\lambda-\mu)^2}, & \lambda>\mu;\end{cases}$$

$$(B-A)(2-|A-B|-A-B)=\begin{cases}\dfrac{\mathcal{D}}{(\lambda-\mu)^2}, & \lambda<\mu,\\[2mm] \dfrac{\mathcal{E}}{(\lambda-\mu)^2}, & \lambda>\mu.\end{cases}$$

Consequently, utilization of [11, Theorem 2.3], [25, Corallary 1] and the first order derivative (16) yields the following conclusions:

(1) The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is increasing on $(0,\infty)$ if and only if $\lambda>\mu$, $\mathcal{A}\ge 0$ and $\mathcal{C}\ge 0$; The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is decreasing on $(0,\infty)$ if and only if $\lambda<\mu$, $\mathcal{A}\ge 0$ and $\mathcal{B}\ge 0$.
(2) The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is decreasing on $(0,\infty)$ if and only if $\lambda>\mu$, $\mathcal{A}\le 0$ and $\mathcal{C}\le 0$; The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is increasing on $(0,\infty)$ if and only if $\lambda<\mu$, $\mathcal{A}\le 0$ and $\mathcal{B}\le 0$.
(3) The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is increasing on $(-\infty,0)$ if and only if $\lambda>\mu$, $\mathcal{A}\ge 0$ and $\mathcal{E}\ge 0$; The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is decreasing on $(-\infty,0)$ if and only if $\lambda<\mu$, $\mathcal{A}\ge 0$ and $\mathcal{D}\ge 0$.
(4) The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is decreasing on $(-\infty,0)$ if and only if $\lambda>\mu$, $\mathcal{A}\le 0$ and $\mathcal{E}\le 0$; The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is increasing on $(-\infty,0)$ if and only if $\lambda<\mu$, $\mathcal{A}\le 0$ and $\mathcal{D}\le 0$.
(5) The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is increasing on $(-\infty,\infty)$ if and only if $\lambda>\mu$, $\mathcal{C}\ge 0$ and $\mathcal{E}\ge 0$; The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is decreasing on $(-\infty,\infty)$ if and only if $\lambda<\mu$, $\mathcal{B}\ge 0$ and $\mathcal{D}\ge 0$.
(6) The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is decreasing on $(-\infty,\infty)$ if and only if $\lambda>\mu$, $\mathcal{C}\le 0$ and $\mathcal{E}\le 0$. The function $H_{\alpha,\beta;\lambda,\mu}(t)$ is increasing on $(-\infty,\infty)$ if and only if $\lambda<\mu$, $\mathcal{B}\le 0$ and $\mathcal{D}\le 0$.

The proof of Theorem 1 is complete. □

Proof of Theorem 2. This follows directly from the combination of Theorem 1 with equations in (8). Theorem 2 is proved. □

References

[1] P. S. Bullen, *Handbook of Means and Their Inequalities*, Mathematics and its Applications (Dordrecht) **560**, Kluwer Academic Publishers, Dordrecht, 2003.
[2] Ch.-P. Chen and F. Qi, *Best constant in an inequality connected with exponential functions*, Octogon Math. Mag. **12** (2004), no. 2, 736–737.
[3] W.-S. Cheung and F. Qi, *Logarithmic convexity of the one-parameter mean values*, RGMIA Res. Rep. Coll. **7** (2004), no. 2, Art. 15, 331–342; Available online at `http://www.staff.vu.edu.au/rgmia/v7n2.asp`.

[4] W.-S. Cheung and F. Qi, *Logarithmic convexity of the one-parameter mean values*, Taiwanese J. Math. **11** (2007), no. 1, 231–237.

[5] P. N. de Souza and J.-N. Silva, *Berkeley Problems in Mathematics*, 2nd ed., Problem Books in Mathematics, Springer, New York, 2001.

[6] N. Elezović, C. Giordano and J. Pečarić, *The best bounds in Gautschi's inequality*, Math. Inequal. Appl. **3** (2000), 239–252.

[7] H. Gauchman, *Steffensen pairs and associated inequalities*, J. Inequal. Appl. **5** (2000), no. 1, 53–61.

[8] B.-N. Guo, A.-Q. Liu and F. Qi, *Monotonicity and logarithmic convexity of three functions involving exponential function*, J. Korea Soc. Math. Educ. Ser. B Pure Appl. Math. **15** (2008), no. 4, 387–392.

[9] B.-N. Guo and F. Qi, *A simple proof of logarithmic convexity of extended mean values*, Numer. Algorithms (2009), in press; Available online at http://dx.doi.org/10.1007/s11075-008-9259-7.

[10] B.-N. Guo and F. Qi, *Generalization of Bernoulli polynomials*, Internat. J. Math. Ed. Sci. Tech. **33** (2002), no. 3, 428–431.

[11] B.-N. Guo and F. Qi, *Properties and applications of a function involving exponential functions*, Commun. Pure Appl. Anal. **8** (2009), no. 4, 1231–1249.

[12] S. Guo and F. Qi, *A class of completely monotonic functions related to the remainder of Binet's formula with applications*, Tamsui Oxf. J. Math. Sci. **25** (2009), no. 1, in press.

[13] J.-Ch. Kuang, *Chángyòng Bùděngshì* (*Applied Inequalities*), 3rd ed., Shandong Science and Technology Press, Ji'nan City, Shandong Province, China, 2004. (Chinese)

[14] A.-Q. Liu, G.-F. Li, B.-N. Guo and F. Qi, *Monotonicity and logarithmic concavity of two functions involving exponential function*, Internat. J. Math. Ed. Sci. Tech. **39** (2008), no. 5, 686–691.

[15] Q.-M. Luo, B.-N. Guo and F. Qi, *Generalizations of Bernoulli's numbers and polynomials*, RGMIA Res. Rep. Coll. **5** (2002), no. 2, Art. 12, 353–359; Available online at http://www.staff.vu.edu.au/rgmia/v5n2.asp.

[16] Q.-M. Luo, B.-N. Guo, F. Qi, and L. Debnath, *Generalizations of Bernoulli numbers and polynomials*, Internat. J. Math. Math. Sci. **2003** (2003), no. 59, 3769–3776.

[17] Q.-M. Luo and F. Qi, *Relationships between generalized Bernoulli numbers and polynomials and generalized Euler numbers and polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **7** (2003), no. 1, 11–18.

[18] F. Qi, *A class of logarithmically completely monotonic functions and the best bounds in the first Kershaw's double inequality*, J. Comput. Appl. Math. **206** (2007), no. 2, 1007–1014.

[19] F. Qi, *A monotonicity result of a function involving the exponential function and an application*, RGMIA Res. Rep. Coll. **7** (2004), no. 3, Art. 16, 507–509; Available online at http://www.staff.vu.edu.au/rgmia/v7n3.asp.

[20] F. Qi, *A note on Schur-convexity of extended mean values*, Rocky Mountain J. Math. **35** (2005), no. 5, 1787–1793.

[21] F. Qi, *Bounds for the ratio of two gamma functions—From Wendel's and related inequalities to logarithmically completely monotonic functions*, Available online at http://arxiv.org/abs/0904.1048.

[22] F. Qi, *Bounds for the ratio of two gamma functions—From Wendel's limit to Elezović-Giordano-Pečarić's theorem*, Available online at http://arxiv.org/abs/0902.2514.

[23] F. Qi, *Logarithmic convexity of extended mean values*, Proc. Amer. Math. Soc. **130** (2002), no. 6, 1787–1796.

[24] F. Qi, *Logarithmic convexities of the extended mean values*, RGMIA Res. Rep. Coll. **2** (1999), no. 5, Art. 5, 643–652; Available online at http://www.staff.vu.edu.au/rgmia/v2n5.asp.

[25] F. Qi, *Monotonicity and logarithmic convexity for a class of elementary functions involving the exponential function*, RGMIA Res. Rep. Coll. **9** (2006), no. 3, Art. 3; Available online at http://www.staff.vu.edu.au/rgmia/v9n3.asp.

[26] F. Qi, *Schur-convexity of the extended mean values*, RGMIA Res. Rep. Coll. **4** (2001), no. 4, Art. 4, 529–533; Available online at http://www.staff.vu.edu.au/rgmia/v4n4.asp.

[27] F. Qi, *The extended mean values: Definition, properties, monotonicities, comparison, convexities, generalizations, and applications*, Cubo Mat. Educ. **5** (2003), no. 3, 63–90.

[28] F. Qi, *Three-log-convexity for a class of elementary functions involving exponential function*, J. Math. Anal. Approx. Theory **1** (2006), no. 2, 100–103.

[29] F. Qi, P. Cerone, S. S. Dragomir and H. M. Srivastava, *Alternative proofs for monotonic and logarithmically convex properties of one-parameter mean values*, Appl. Math. Comput. **208** (2009), no. 1, 129–133; Available online at http://dx.doi.org/10.1016/j.amc.2008.11.023.

[30] F. Qi and J.-X. Cheng, *New Steffensen pairs*, RGMIA Res. Rep. Coll. **3** (2000), no. 3, Art. 11, 431–436; Available online at http://www.staff.vu.edu.au/rgmia/v3n3.asp.

[31] F. Qi and J.-X. Cheng, *Some new Steffensen pairs*, Anal. Math. **29** (2003), no. 3, 219–226.

[32] F. Qi, J.-X. Cheng and G. Wang, *New Steffensen pairs*, Inequality Theory and Applications, Volume **1**, 273–279, Nova Science Publishers, Huntington, NY, 2001.

[33] F. Qi and B.-N. Guo, *An alternative proof of Elezović-Giordano-Pečarić's theorem*, Available online at http://arxiv.org/abs/0903.1174.

[34] F. Qi and B.-N. Guo, *Generalisation of Bernoulli polynomials*, RGMIA Res. Rep. Coll. **4** (2001), no. 4, Art. 10, 691–695; Available online at http://www.staff.vu.edu.au/rgmia/v4n4.asp.

[35] F. Qi and B.-N. Guo, *On Steffensen pairs*, J. Math. Anal. Appl. **271** (2002), no. 2, 534–541.

[36] F. Qi and B.-N. Guo, *On Steffensen pairs*, RGMIA Res. Rep. Coll. **3** (2000), no. 3, Art. 10, 425–430; Available online at http://www.staff.vu.edu.au/rgmia/v3n3.asp.

[37] F. Qi and B.-N. Guo, *The function* $(b^x - a^x)/x$*: Logarithmic convexity*, RGMIA Res. Rep. Coll. **11** (2008), no. 1, Art. 5; Available online at http://www.staff.vu.edu.au/rgmia/v11n1.asp.

[38] F. Qi and B.-N. Guo, *The function* $(b^x - a^x)/x$*: Logarithmic convexity and applications to extended mean values*, Available online at http://arxiv.org/abs/0903.1203.

[39] F. Qi and B.-N. Guo, *Wendel's and Gautschi's inequalities: Refinements, extensions, and a class of logarithmically completely monotonic functions*, Appl. Math. Comput. **205** (2008), no. 1, 281–290; Available online at http://dx.doi.org/10.1016/j.amc.2008.07.005.

[40] F. Qi and B.-N. Guo, *Wendel-Gautschi-Kershaw's inequalities and sufficient and necessary conditions that a class of functions involving ratio of gamma functions are logarithmically completely monotonic*, RGMIA Res. Rep. Coll. **10** (2007), no. 1, Art. 2; Available online at http://www.staff.vu.edu.au/rgmia/v10n1.asp.

[41] F. Qi, B.-N. Guo and Ch.-P. Chen, *The best bounds in Gautschi-Kershaw inequalities*, Math. Inequal. Appl. **9** (2006), no. 3, 427–436.

[42] F. Qi, S. Guo and B.-N. Guo, *A class of* k*-log-convex functions and their applications to some special functions*, RGMIA Res. Rep. Coll. **10** (2007), no. 1, Art. 21; Available online at http://www.staff.vu.edu.au/rgmia/v10n1.asp.

[43] F. Qi, S. Guo, B.-N. Guo and Sh.-X. Chen, *A class of* k*-log-convex functions and their applications to some special functions*, Integral Transforms Spec. Funct. **19** (2008), no. 3, 195–200.

[44] F. Qi and Q.-M. Luo, *A simple proof of monotonicity for extended mean values*, J. Math. Anal. Appl. **224** (1998), no. 2, 356–359.

[45] F. Qi, D.-W. Niu and B.-N. Guo, *Monotonic properties of differences for remainders of psi function*, Internat. J. Pure Appl. Math. Sci. **4** (2007), no. 1, ???–???.

[46] F. Qi, D.-W. Niu and B.-N. Guo, *Monotonic properties of differences for remainders of psi function*, RGMIA Res. Rep. Coll. **8** (2005), no. 4, Art. 16, 683–690; Available online at http://www.staff.vu.edu.au/rgmia/v8n4.asp.

[47] F. Qi and S.-L. Xu, *Refinements and extensions of an inequality, II*, J. Math. Anal. Appl. **211** (1997), no. 2, 616–620.

[48] F. Qi and S.-L. Xu, *The function* $(b^x - a^x)/x$*: Inequalities and properties*, Proc. Amer. Math. Soc. **126** (1998), no. 11, 3355–3359.

[49] F. Qi, S.-L. Xu and L. Debnath, *A new proof of monotonicity for extended mean values*, Internat. J. Math. Math. Sci. **22** (1999), no. 2, 415–420.

[50] Y.-D. Wu and Zh.-H. Zhang, *The best constant for an inequality*, Octogon Math. Mag. **12** (2004), no. 1, 139–141.

[51] Y.-D. Wu and Zh.-H. Zhang, *The best constant for an inequality*, RGMIA Res. Rep. Coll. **7** (2004), no. 1, Art. 19; Available online at http://www.staff.vu.edu.au/rgmia/v7n1.asp.

[52] Sh.-Q. Zhang, B.-N. Guo and F. Qi, *A concise proof for properties of three functions involving the exponential function*, Appl. Math. E-Notes **9** (2009), in press.

(B.-N. Guo) School of Mathematics and Informatics, Henan Polytechnic University, Jiaozuo City, Henan Province, 454010, China
E-mail address: bai.ni.guo@gmail.com, bai.ni.guo@hotmail.com

(F. Qi) Research Institute of Mathematical Inequality Theory, Henan Polytechnic University, Jiaozuo City, Henan Province, 454010, China
E-mail address: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com
URL: http://qifeng618.spaces.live.com